arXiv:1304.2887v2 [quant-ph] 18 Apr 2013

# Non classicality of generalized vortex states produced by multi-photon subtraction from two mode squeezed vacuum

Anindya Banerji[1]*, Ravindra Pratap Singh[2]†, and Abir Bandyopadhyay[1]‡
[1] *Hooghly Engineering and Technology College, Hooghly 712103, India and*
[2] *Physical Research Laboratory, Ahmedabad 380009, India*

(Dated: April 19, 2013)

We discuss the non classicality of a class of vortex states which are engineered by photon substraction from two mode squeezed vacuum. We calculate the log negativity parameter, the von Neumann entropy of the reduced density matrix and the concurrence. The negativity of the Wigner function is discussed as a measure of the non classicality and the quantum interference pattern obtained therein as a possible measure of the entanglement present between the two modes. This measure of entanglement is compared with the results obtained from concurrence.

PACS numbers: 42.65.Lm, 03.65.Ud, 03.67.Mn, 03.67.Bg, 89.70.Cf
Keywords: Quantum optical vortex, Photon addition/subtraction, Wigner function, Entanglement, Entropy

## I. INTRODUCTION

Over the last decade, the physical characteristics of the entanglement of quantum mechanical states, both pure and mixed, has been recognised as a central resource in various aspects of quantum information processing [1]. Given an arbitrary bipartite pure state $|\Psi_{AB}\rangle$, the entropy of entanglement $E(|\Psi_{AB}\rangle)$ [2], namely the von Neumann entropy of the reduced density matrix $\rho_A \equiv \mathrm{Tr}_B|\Psi_{AB}\rangle\langle\Psi_{AB}|$, tells us exhaustively about the probabilities of transforming $|\Psi_{AB}\rangle$ using LOCC, into other pure states in an asymptotic sense. Many efforts have also been devoted to the study of mixed state entanglement [3, 4]. Given the central status of entanglement, the task of quantifying the degree to which a state is entangled is important for quantum information processing and correspondingly several measures such as entanglement of formation [3, 5], von Neumann entropy [6] and negativity [1, 7] have been formulated. The entanglement of formation or concurrence have been the topic of much interest lately. Although originally formulated to quantify pure state entanglement, it was later extended to mixed states [8–11].

Among all (quasi) probability distributions in quantum mechanics [12], Wigner function [13] stands out because it is real, non singular, yields correct quantum mechanical operator averages in terms of phase space integrals and possesses definite marginal distribution [14]. Moreover, the negativity of Wigner function depicts non classicality of the state [15], which we highlight in this article as a measure of entanglement also.

Optical vortices or phase singularities in the field of light, described by a non-separable two-dimensional field, have drawn a great deal of attention in the last two decades [16] and these have prompted to start a new branch in physical optics known as singular optics [17]. Because of their specific spatial structure and associated orbital angular momentum (OAM), they find various applications in the field of optical manipulation [18], optical communication [19], quantum information and computation [20]. For a vortex, the OAM follows the same numbers as vorticity or topological charge of the vortex i.e. an optical vortex of topological charge $m$, carries an OAM of $m\hbar$ per photon. Most applications of vortices lie in the classical realm. However, in recent literature [21–25] quantum optical vortex has also been studied. It has been shown that such states can be generated from two mode squeezed vacuum under a linear transformation belonging to the SU(2) group with some restrictions.

We studied the Wigner function of a general elliptical vortex, the circular vortex being a special case of an elliptical vortex which is constructed from two separate squeezed vacuum states by mixing them through a beam splitter (BS) or a dual channel directional coupler (DCDC) [23]. We also studied the general elliptical vortex from an information theoretic viewpoint [26]. We looked at the suitability of it for carrying information by studying the von Neumann entropy.

Recently Agarwal [24] analyzed quantum optical vortex by subtracting a photon by a 99% transmitting beam - splitter (BS) from one of the two modes of spontaneous parametric down-conversion (SPDC), generated by the non - linear crystal (NLC). He showed that the subtraction from one mode results into an addition to the other mode; it produces a vortex, which is elliptical in shape. It must be noted that a single photon subtraction or addition has already been demonstrated by Parigi et al. [27]. A more recent paper studied the enhancement of entanglement, though defined differently, in photon subtraction/addition [28]

The paper is organized as follows. In section II, we introduce the generalized vortex state arising from multi - photon subtraction of the two mode squeezed vacuum. We show that multi - photon subtraction indeed gives rise to vortex states of higher order. We discuss the non classicality of this state with the help of the Wigner function.

---

*abanerji09@gmail.com
†rpsingh@prl.res.in
‡abir@hetc.ac.in

In section III, we discuss the entanglement properties of the generalized vortex state with the logarithmic negativity and entanglement of formation. We discuss the variation in mixedness with squeezing. We also present an interpretation of the quantum interference patterns obtained from the Wigner function as a measure of the entanglement present between the two modes by comparing it with the results obtained from our study of concurrence. We conclude this article after pointing out the significant results and directions for future work in section IV.

## II. VORTEX STATES ARISING FROM PHOTON SUBTRACTION/ADDITION

Quantum optical vortices are phase space singularities in the quantized radiation field which possess orbital angular momentum. There are methods of producing these states. One of them is by mixing two squeezed vacuum modes through BS or dual channel directional coupler [23]. The other one is by photon subtraction using a BS from one of the outputs of a SPDC [24, 27] as shown in Fig. 1. Vortex of higher orders can be produced by increasing the number of BSs and avalanche photo diodes. Here we discuss the properties of vortices created by subtracting photons.



FIG. 1: A schematic of the experimental setup for the production of vortex state of order 2 by subtracting two photons from a two - mode squeezed vacuum using two 99% beam splitters(BS 1 and BS 2). NLC is a nonlinear crystal. SPDC generates the signal (mode a) and the idler (mode b). APD 1 and 2 are Avalanche Photo Diodes.

A two-mode squeezed beam is represented by

$$|\xi\rangle = \exp\left(\xi a^{\dagger}b^{\dagger} - \xi^{*}ab\right)|0,0\rangle, \quad \xi = re^{i\phi} \tag{1}$$

where $\xi$ is a complex parameter, $r$ is the squeezing parameter and $a$ and $b$ are the regular bosonic mode operators satisfying the commutation relations $[a, a^{\dagger}] = [b, b^{\dagger}] = 1, [a, b^{\dagger}] = 0$, etc. The corresponding quadrature distribution is obtained by replacing the bosonic mode operators by the quadrature operators $x, y, p_x$ and $p_y$ which obey the relations $[x, p_x] = [y, p_y] = i$. Here we consider $\hbar = 1$. The quadratures $x$, $p_x$ correspond to mode $a$ (signal) and $y$, $p_y$ correspond to mode $b$ (idler). The quadrature distribution thus obtained has the following form

$$\Psi^{(s)}(x,y) = \frac{1}{\sqrt{\left(1-\eta^{2}\right)\pi\cosh^{2}r}} \times\exp\left[\frac{2xy\eta - \left(x^{2}+y^{2}\right)\eta^{2}}{1-\eta^{2}} - \frac{1}{2}\left(x^{2}+y^{2}\right)\right] \tag{2}$$

The non-separability is evident from the fact that Eq.



(a)

(b)

(c)

FIG. 2: (Color online) Plot of the intensity (a) 3D plot, (b) contour plot and (c) phase of $\Psi_k^s(x,y)$ for $k$=3. The singularities are clearly seen in Fig. (c).

(2) cannot be written as a product of two functions each of which are solely dependent on either $x$ or $y$.
Now one photon can be subtracted from mode $b$ with the help of a beam splitter.

$$|\xi\rangle^{(s)} = \mathcal{A} b \exp\left(\xi a^\dagger b^\dagger - \xi^* ab\right)|0,0\rangle \tag{3}$$

where $\mathcal{A}$ is the constant of normalization. This photon can be detected using a single photon detector. The probability of detection is proportional to the square of reflectivity [24]. If an arbitrary number of photons, say $k$, are subtracted from the two mode squeezed vacuum, Eq. (3) can be generalized to the following

$$|\xi\rangle_k^{(s)} = \mathcal{A}_k b^k \exp\left(\xi a^\dagger b^\dagger - \xi^* ab\right)|0,0\rangle \tag{4}$$

We can rewrite Eq. (4) as follows

$$\begin{aligned} |\xi\rangle_k^{(s)} &= \mathcal{A}_k b^{k-1} \exp\left(\xi a^\dagger b^\dagger - \xi^* ab\right) \\ &\times \exp\left(-\xi a^\dagger b^\dagger + \xi^* ab\right) b \exp\left(\xi a^\dagger b^\dagger - \xi^* ab\right)|0,0\rangle \end{aligned} \tag{5}$$

This can be simplified by repeated use of the two mode Bogoliubov transformation to finally arrive at the following simplified form

$$|\xi\rangle_k^{(s)} = \mathcal{A}_k \exp\left(\xi a^\dagger b^\dagger - \xi^* ab\right)\left(b\mathrm{cosh}r + a^\dagger \mathrm{sinh}r e^{i\theta}\right)^k |0,0\rangle \tag{6}$$

Eq. (6) can be further simplified to

$$|\xi\rangle_k^{(s)} = \mathcal{A}_k \exp\left(\xi a^\dagger b^\dagger - \xi^* ab\right)|k,0\rangle \sinh^k r e^{ik\theta} \tag{7}$$

One can also write the state (Eq. (7)) in terms of Fock states by using the decomposition of the squeezing operator and obtain

$$\begin{aligned} |\xi\rangle_k^{(s)} &= \frac{e^{ik\theta}}{\cosh^2 r}\sum_{m=0}^{\infty} e^{im\theta}(\tanh^m r\sqrt{m+k}|m+k,m\rangle \\ &= \frac{e^{ik\theta}}{\cosh^2 r} a^{\dagger k}|\xi\rangle \end{aligned} \tag{8}$$

So we see that the state (8) can also be obtained by adding $k$ photons to $|\xi\rangle$ in the mode $a$. This is similar to Parigi *et al*'s experiment in which they added a single photon to generate the vortex state. The state (8) exhibits the property that there is a fixed difference in the number of photons between the signal and idler states. In literature this is one of the so called "pair coherent states" [29, 30]. Pair coherent states provide an important example of non classical states of the two mode radiation field. It has been studied in detail for their non classical properties and as examples of EPR states [29].

The other property of interest is the vortex structure. The quadrature distribution for the state (8) can be obtained by writing the bosonic creation operator $a^\dagger$ in terms of $x_a$ and $\partial/\partial x_a$ as follows

$$\Psi_k^{(s)}(x,y) = \frac{e^{ik\theta}}{2^{k/2}\mathrm{cosh r}}\left(x - \frac{\partial}{\partial x}\right)^k \Psi^{(s)}(x,y) \tag{9}$$



(a)

(b)

(c)

FIG. 3: (Color online) Plot of the intensity (a) 3D plot, (b) contour plot and (c) phase of $\Psi_k^s(x,y)$ for $k$=4. The singularities are clearly seen in Fig. (c).

where $\Psi^{(s)}(x,y)$ is the quadrature distribution of the two mode squeezed vacuum, Eq. (2). For $k=1$, Eq. (9) can be expressed as follows

$$\begin{aligned} \Psi_1^{(s)}(x,y) &= \frac{\sqrt{2}e^{i\theta}\left(x-\eta y\right)}{\left(1-\eta^2\right)^{3/2}\pi^{1/2}\cosh^2 r} \\ &\times \exp\left[\frac{2xy\eta - \left(x^2+y^2\right)\eta^2}{1-\eta^2} - \frac{1}{2}\left(x^2+y^2\right)\right] \end{aligned} \tag{10}$$

This is seen to possess a vortex structure for $\eta = i|\eta|$. In other words, the method of single photon subtraction



(a) $W(x,y)^{p_x=0}_{p_y=0}$ (b) $W(p_x,p_y)^{x=0}_{y=0}$ (c) $W(x,p_x)^{y=0}_{p_y=0}$

(d) $W(y,p_y)^{x=0}_{p_x=0}$ (e) $W(x,p_y)^{p_x=0}_{y=0}$ (f) $W(y,p_x)^{x=0}_{p_y=0}$

FIG. 4: Wigner function of the TMSV state, for the five photon subtracted case (k=5).

from one of the two modes taken from the output of a spontaneous parametric downconverted beam results in the formation of a vortex state of order 1. Intuitively it can be argued that the process of subtracting $k$ photons should give rise to vortex states of higher order. As has already been pointed out, multi-photon subtraction from one of the modes of the two mode squeezed vacuum is similar to adding an equal number of photons in the other mode. The basic idea is to have a difference in the photon number between the two modes. This gives rise to higher order vortex states where the order or vorticity is determined by the difference in number of photons in the two modes. We call these states Two Mode Squeezed Vortex (TMSV) states.

We study the states arising from 3 photon and 4 photon subtraction in Fig. (2) and Fig. (3) respectively. It is evident that this method gives rise to higher order vortex states. The order can be determined from the phase plots by counting the number of singular points. For $k$=1, the results of [24] is reproduced. It is observed, with increasing order, the intensity distribution associated with the vortex begins to divide into separate lobes, as can be seen from Fig. (2), when three photons are subtracted ($k$=3). When another photon is subtracted, the corresponding intensity distribution is completely divided into two distinct lobes. These results serve as a pointer to study the entanglement properties of these states and its variation with the order of the vortex.

The non classical nature of these states can be studied with the help of the Wigner function. For a number state $|n\rangle$, it is written as [31],

$$W(x,y) = \frac{2}{\pi}\,(-1)^n\,\mathcal{L}_n\left(4\,\boldsymbol{r}^2\right)e^{-2\,\boldsymbol{r}^2}, \tag{11}$$

where $\boldsymbol{r}^2 = x^2 + y^2$ and $\mathcal{L}_n$ is the Laguerre polynomial of order $n$. The Wigner function for a two mode Fock state, $|n,m\rangle$, can then be obtained in terms of coherent state parameters $\alpha = x - ip_x$ and $\beta = y - ip_y$ as follows

$$\begin{aligned} W(\alpha,\beta) &= \frac{4}{\pi^2}\,(-1)^n\,\mathcal{L}_n\left[4|\alpha|^2\right]\exp\left(-2|\alpha|^2\right) \\ &\times\ (-1)^m\,\mathcal{L}_m\left[4|\beta|^2\right]\exp\left(-2|\beta|^2\right) \end{aligned} \tag{12}$$

The Wigner function for the two mode state $|n,0\rangle$ can then be simplified and written as

$$W(\alpha,\beta) = \frac{4}{\pi^2}\,(-1)^n\,\mathcal{L}_n\left[4|\alpha|^2\right]\exp\left[-2\left(|\alpha|^2+|\beta|^2\right)\right]. \tag{13}$$

It has been shown that if two density matrices are related by a squeezing transformation, then the Wigner functions are related by the following transformation [24]

$$W(\alpha,\beta) = W(\tilde{\alpha},\tilde{\beta}), \tag{14}$$

where,

$$\begin{pmatrix}\tilde{\alpha}\\ \tilde{\beta}^*\end{pmatrix} = \begin{pmatrix}\cosh r & -\sinh r\ e^{i\theta}\\ -\sinh r\ e^{i\theta} & \cosh r\end{pmatrix}\begin{pmatrix}\alpha\\ \beta^*\end{pmatrix} \tag{15}$$

So, the Wigner function for the TMSV, $|\xi\rangle^{(s)}_k$, can be written as,

$$W(\tilde{\alpha},\tilde{\beta}) = \frac{4}{\pi^2}\,(-1)^k\,\mathcal{L}_k\left[4|\tilde{\alpha}|^2\right]\exp\left[-2\left(|\tilde{\alpha}|^2+|\tilde{\beta}|^2\right)\right], \tag{16}$$

where $k$ is the number of photons subtracted. In Fig. (4) we study the variation of the Wigner function as a function of two of the quadratures out of four (two quadratures for each of the two modes), keeping the other two quadratures as zero.

The non classicality of the TMSV states is evident from the nature of the Wigner function which shows negative regions for all the photon subtracted states. In Fig. (4a) and Fig. (4b), we study the cross correlation between the same quadratures of two different modes. It is observed, though not shown in the figure, that the peak alternates between 0.4 (for even $k$) and -0.4 (for odd $k$). It attains the positive maximum when an even number of photons are subtracted. Negative maximum is attained when an odd number of photons are subtracted. The volume of the negative region of the Wigner functions can be further studied as a measure of the non classicality of these states [15]. It can also be interpreted as a measure of entanglement between the two modes as explained in section III C. Fig. (4c) and Fig. (4d) represents the cross correlation between two quadratures of the same mode. These are exactly similar in nature to that of the cross correlation between same quadratures of different modes. In Fig. (4e) and Fig. (4f), we study the cross correlation between different quadratures of two different modes. These show interesting quantum interference effects. Similar interference pattern has been reported in [22, 32]. We present an interpretation of this quantum inteference later in this article.

## III. ENTANGLEMENT PROPERTIES

In this section we study the mixedness of the TMSV states as well as the entanglement properties of them.

### A. Logarithmic negativity

There exists various measures of entanglement. A much used tool is the log negativity parameter [1]. We use this to quantatively study the entanglement in the vortex state. It is defined by

$$\epsilon = \log_2 (1 + 2\mathcal{N}) \tag{17}$$

where $\mathcal{N}$ is the modulus of the sum of all the negative eigenvalues associated with the partial transpose $\rho^{\mathrm{PT}}$ of the density matrix $|\xi\rangle_k^{(s)}\,{}_k^{(s)}\langle\xi|$. The density matrix of the TMSV state can be written from Eq. (8) as follows

$$\rho = \sum_{m,n} c_m c_n |m+k, m\rangle\langle n+k, n| e^{i(m-n)\theta} \tag{18}$$

where,

$$c_p = \tanh^p r \frac{\sqrt{p+1}}{\cosh^2 r}. \tag{19}$$

Here $p = m,\ n$. Therefore, the log negativity parameter of Eq. (17) becomes

$$\epsilon_k = \log_2 \left(\sum_n c_n^2\right)^2 . \tag{20}$$



(a) Two photon subtraction



(b) Three photon subtraction

FIG. 5: Entanglement ratio $\tilde{\epsilon_k}$ for the TMSV states

A similar calculation for the two mode squeezed vacuum gives the result $\log_2\left(e^{2r}\right)$ using $c_n = \tanh^n r/\cosh r$. The entanglement in the vortex state can be compared with that in the state $|\xi\rangle$ by studying the ratio of log negativities of the two states. The ratio has the form

$$\tilde{\epsilon}_k = \left(\sum_n c_n e^{-r}\right)^2 . \tag{21}$$

We study the results in Fig. (5). For $k = 1$, i.e. single photon subtraction, the results of [24] were reproduced. It was observed that for two photon subtraction there was a very small rise in entanglement which falls sharply after reaching the peak. Further subtraction worsens the situation. For three photon subtraction the initial value falls to half. For four photon subtraction, the starting value is 0.042. This seems pretty interesting given the nature of the higher order vortex states in Fig. (2) and Fig. (3). When three photons are subtracted, it gives rise to a vortex state of order 3 as can be seen from Fig. (2c). But it is also evident from Fig. (2b) that the corresponding intensity distribution almost disintegrates into separate lobes. The situation worsens when four photons are subtracted (Fig. (3c) and Fig. (3b)). The intensity distribution, in this case, is seen to have completely divided into two distinctly separate lobes. Although the lobes are separate, it does not imply that there is no entanglement between photons sampled from the different lobes.

### B. von Neumann entropy

The von Neumann entropy, also known as the entropy of entanglement is defined as

$$S_a = -\mathrm{Tr}\left(\rho_a \log_2 \rho_a\right) = -\sum_n \lambda_n \log_2 \lambda_n, \qquad (22)$$

where, $\lambda_n$ are the eigenvalues of $\rho_a$, the reduced density matrix of $|\xi\rangle_k^{(s)}$. The reduced density matrix is defined as $\mathrm{Tr}_a\left[|\xi\rangle_k^{(s)}\ {}_k^{(s)}\langle\xi|\right]$. Using Eq. (18), we can write it down as follows

$$\rho_a = \sum_k c_k^2 |m+1\rangle\langle m+1|, \qquad (23)$$

where $c_k$,s are the same coefficients as in Eq. (19). The entropy of entanglement, as is well known, can be interpreted as a measure of the mixedness of the states. Its values, when appropriately scaled, vary between zero, for pure states, and one, for completely mixed states. We study the variation of the same with the squeezing parameter, in Fig. (6).



FIG. 6: Entropy of entanglement for the TMSV states for $k$=4.

We observe, that the entropy increases with the squeezing parameter, $r$. It increases from 0 to reach a maximum value, at $r = 3.2$. Since von Neumann entropy is interpreted as a measure of the mixedness [10], this implies that the states change from completely pure to completely mixed state with the increase in squeezing. When $r = 0$, that is, there is no squeezing, the states remain pure. It gradually looses its purity with increasing $r$ and becomes completely mixed at a higher value of $r$. The dependence of the mixedness of the TMSV on the squeezing parameter is not very surprising. In our setup, we use the SPDC to squeeze the two mode vacuum. This also results in the formation of a mixed state. So, the squeezing parameter apart from parametrizing the squeezing of the two mode vacuum also parametrizes the mixing of the two modes. It is also observed that the variation of the von Neumann entropy with squeezing parameter is independent of the order of the vortex.

### C. Entanglement of formation

In this section we study the entanglement of the TMSV state with the help of concurrence [5, 9], also expressed as the entanglement of formation [3]. It is a measure of the amount of entanglement needed to create the entangled state. It is defined as [10]

$$C(\rho) = \min_{p_i,\psi_i} \sum_i p_i E\left(|\psi_i\rangle\langle\psi_i|\right) \qquad (24)$$

The minimization is taken over all possible decompositions of pure states $\psi_i$ with probabilities $p_i$, which taken together reproduce the density matrix $\rho$ of the given mixed state. On the other hand, $E\left(|\psi_i\rangle\langle\psi_i|\right)$ is the entropy of entanglement, more commonly known as the von Neumann entropy, of the reduced density matrix as defined in the previous section III B. Using Eq. (18), Eq. (23) and Eq. (22), we arrive at the following simplified form of the entanglement of formation



FIG. 7: (Color Online) Entanglement of formation of the photon added (subtracted) vortex state. $m$ denotes the number of photons added (subtracted) in one of the modes.

$$C(\rho)_k = \min_{c_n, |\xi\rangle_k^{(s)}} \sum_{n,m} c_n c_m \left(-\sum_p c_p^2 \log_2 c_p^2\right) \qquad (25)$$

where, $c_i$'s are the same as in Eq. (19). We study the variation of the entanglement of formation with the squeezing parameter and the order of the vortex in Fig. (7). We observe that the $C(\rho)_k$ increases with increase in $r$, the squeezing parameter. The rate of increase gradually slows down till it reaches the maximum value and then falls of rapidly to 0. It should be noticed that $C(\rho)_k$ attains its maximum value at $r \sim 2.1$. This means that the two modes become more and more entangled with increasing $r$ until it reaches a maximum value beyond which the modes start becoming disentangled. We infer that there exists an optimum value of the squeezing parameter which when applied at the beam splitter produces maximum entanglement of the two modes. If we go on squeezing the modes more and more, they will no more be entangled. As we move to higher orders of the vortex, the range of values of $r$ which produces maximal entanglement, starts broadening. At $k$=5, there exists



(a) r=0.4



(b) r=1.5



(c) r=2.1



(d) r=3

FIG. 8: Wigner function $W(x, p_y)$ of the TMSV states showing quantum interference for different values of $r$, the squeezing parameter, with $k = 5$.

a well defined range of $r$ that produces maximal entanglement of the two modes. The entanglement falls off rapidly on both the ends of this range.

We present here an interpretation of the quantum interference effects observed in section II. In Fig. (8), we study the effect of the squeezing parameter on quantum interference. It is observed that the interference is most prominent for $r = 2.1$ (see Fig. 8c). This observation has a striking similarity with the results obtained from our study of the entanglement of formation. The interference effects decrease as we move away from $r = 2.1$ on either side which is exactly similar to what we obtained from the entanglement of formation. Broadening of the range of $r$ which gives rise to a well defined interference fringe system with increase in the order of the vortex was also observed. Hence, we suggest that the quantum interefence effects arising from the Wigner function of the cross correlation between different quadratures of the two modes can be interpreted as a signature of the entanglement present between the two modes. It is seen that the number of fringes as well as the volume of the fringes change with changing $r$. However, the maximum number of negative fringes is restricted by the order of the vortex. Since the negativity of the Wigner function is interpreted as a measure of non classicality, we suggest, the volume of the negative part of the quantum interference fringes might be a possible candidate for quantifying entanglement and needs to be further studied.

## IV. CONCLUSION

In conclusion, we have studied the non classical properties of the photon subtracted (added) two mode squeezed vacuum state. We have shown that the order of the vortex created by photon subtraction (addition) is equal to the number of photons subtracted (added) from one of the modes. Rather it should be stated that the order was equal to the difference in the number of photons between the two modes achieved by subtracting (adding) photons from one of the two modes. The intensity distribution of the higher order vortex, thus created, starts disintegrating. This gives rise to two distinctly separate lobes.

The non classicality was evident from the negative regions of the Wigner function. The volume of the negative region can be interpreted as a measure of the non classicality and should be a topic of further investigation. We studied the mixedness of these states with the help of von Neumann entropy and the entanglement properties using the logarithmic negativity and the entanglement of formation. The states were observed to change from pure to mixed states with increase in squeezing. It was found that there exists a threshold value of the squeezing parameter, beyond which the states become completely mixed. It was observed that the entanglement decreases with increasing order of the vortex with the help of the logarithmic negativity. However, no such evidence was found while studying the entanglement of formation or concurrence. On the contrary, it was observed that with increasing order of the vortex, the range of values of $r$, the squeezing parameter, which gives rise to optimum entanglement, broadened. It was found from the study of the entanglement of formation that the maximum entanglement achieved for all orders of the vortex was equal. The only difference was the range of values of $r$ for which it was possible to achieve the maximum entanglement. The higher the order of the vortex, the broader was the range of $r$.

It is also suggested that the quantum interference effects obtained from the study of the cross correlation between different quadratures of two different modes might be interpreted as arising due to entanglement between the two modes. The volume of the negative regions of the interference fringes need to be further investigated as a possible measure of the entanglement. This further points to the need of a more robust measure of entanglement.

The present measures of entanglement are found to be inadequate in that they are dependent on the system of states being studied and are not universal measures. The TMSV state discussed in this article should also be further investigated to see how it behaves under teleportation.

### Acknowledgments

This work is partially supported by DST through SERB grant no.: SR/S2/LOP - 0002/2011. A Bandyopadhyay acknowledges the Associateship at Physical Research Laboratory, Ahmedabad, where part of the work is done.

---

[1] G. Vidal, R. F. Werner, Phys. Rev. A **65** (2002) 032314.
[2] C. H. Bennett *et al*, Phys. Rev. A **53** (1996) 2046.
[3] C. H. Bennett *et al*, Phys. Rev. A **54** (1996) 3824; arXiv:quant-ph/9604024.
[4] M. Horodecki, P. Horodecki, R. Horodecki, Phys. Rev. Lett **80** (1998) 5239; arXiv:quant-ph/9801069; *Quantum Information*, Springer (Berlin Heidelberg), 2001; arXiv:quant-ph/0109124.
[5] W. Wootters, Phys. Rev. Lett. **80** (1998) 2245; arXiv:quant-ph/9709029.
[6] M. A. Nielson, I. L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, 2000.
[7] Zyczkowski *et al*, Phys. Rev. A **58** (1998) 883.
[8] W. Munro, J. Opt. B **2** (2000) 47.
[9] W. Wootters, *Quantum Information and Computation*, **1** (2001) 27.
[10] Tzu-Chich Wei *et al*, Phys. Rev. A **67** (2003) 022110.
[11] K. Chen, S. Albeverio, Shao-Ming Fei, arXiv:quant-ph/0506136v2 (2005).
[12] K. Husimi, Proc. Phys. Math. Soc. Jpn. **22** 264 (1940); R. J. Glauber, Phys. Rev. **131** 2766 (1963); E. C. G. Sudarshan, Phys. Rev. Lett. **10** 277 (1963).
[13] E. P. Wigner, Phys. Rev. **40** 749 (1932).
[14] M. Hillery *et al*, Phys. Rep. **106** (1984) 121; H. W. Lee, Phys, Rep. **259** (1995) 147; W. P. Schleich, *Quantum Optics in Phase Space*, Wiley-VCH, Berlin, 2001.
[15] A. Kenfack, K. Życzkowski, J. Opt. B: Q. Semiclass. Opt. **6** (2004) 396.
[16] L. Allen *et al*, *Progress in Optics*, **39** ed. E. Wolf (1999) 291; J. Leach *et al*, Phys. Rev. Lett. **88** (2002) 257901; R. P. Singh, S. Roychowdhury, V. K. Jaiswal, J. Mod. Opt. **53**, (2006) 1803.
[17] M. S. Soskin, M. V. Vasnetsov, *Progress in Optics* **41** ed. E. Wolf (2000) 219.
[18] D. G. Grier, Nature **424** (2003) 810.
[19] G. Gibson *et al*, Opt. Exp. **12** (2004) 5448.
[20] A. Mair *et al*, Nature **412** (2001) 313.
[21] G. S. Agarwal, R. R. Puri, R. P. Singh, Phys. Rev. A **56** (1997) 4207.
[22] G. S. Agarwal, J. Banerji, J. Phys. A **39** (2006) 11503.
[23] A. Bandyopadhyay, R. P. Singh, Opt. Comm. **284** (2011) 256.
[24] G. S. Agarwal, New J. Phys. **13** (2011) 073008.
[25] A. K. Jha, G. S. Agarwal, R. W. Boyd, Phys. Rev. A **83** (2011) 053829.
[26] A. Banerji, P. K. Panigrahi, R.P. Singh, S. Chowdhury, A. Bandyopadhyay, arXiv:quant-ph/1210.6744 (2012).
[27] V. Parigi, A. Zavetta, M. Kim, M. Bellini, Science **317** (2007) 1890.
[28] C. Nvarette-Benlloch *et al*, Phys. Rev. A **86** (2012) 012328.
[29] G. S. Agarwal, Phys. Rev. Lett. **57** (1986) 827; J. Opt. Soc. Am. B **5** (1988) 1940; K. Tara, G. S. Agarwal, Phys. Rev. A **50** (1994) 2870; G. S. Agarwal, A. Biswas, J. Opt. B: Quant. Semiclass. Opt, **7**, (2005) 350.
[30] Arvind, Phys. Lett. A **299** (2002) 461.
[31] D. F. Walls, G. J. Milburn *Quantum Optics*, Springer - Verlag (Berlin Heidelberg), 1994.
[32] S. A. Ramakrishna, A. Bandyopadhyay, J. Rai, Opt. Exp. **2** (1998) 29.